
ASSESS COLD PATIENT

- 1. From outside ring to centre: assess Consciousness, Movement, Shivering, Alertness
- 2. Assess whether normal function, or impaired or no function
- 3. Treat according to appropriate result-quadrant
- 4. Treat all traumatized cold patients with active warming to upper trunk



CARE FOR COLD PATIENT

SUGGESTED SUPPLIES FOR SEARCH/RESPONSE TEAMS IN COLD ENVIRONMENTS:

- 1 Tarp or plastic sheet for vapour barrier outside sleeping bag
- 1 Plastic or foil sheet (2 x 3 m) for vapour barrier placed inside sleeping bag
- 1 Insulated ground pad
- 1 Hooded sleeping bag (or equivalent)
- 1 Source of heat for each team member (e.g., chemical heating pads, or warm water in a bottle or hydration bladder), or each team (e.g., charcoal heater, chemical / electrical heating blanket, or military style Hypothermia Prevention and Management Kit [HPMK])

INSTRUCTIONS FOR HYPOTHERMIA WRAP "The Burrito"


1. Dry or damp clothing: Leave clothing on

2. Verv wet clothina:

THEN Wrap immediately IF Shelter / Transport is more than 30 minutes away, THEN Protect patient from environment,

IF Shelter / Transport is less than 30 minutes away.

remove wet clothing and wrap



Copyright © 2016. Baby It's Cold Outside. All rights reserved. BICOrescue.com Sources; BICOrescue.com; Zafren, Giesbrecht, Danzl et al, Wilderness Environ Med, 2014, 25:S66-85.